

Leibniz Homology, Characteristic Classes and K-theory

Jerry M. Lodder*

1 Introduction

In this paper we identify many striking elements in Leibniz (co)homology which arise from characteristic classes and K-theory. For a group G and a field k containing $\mathbf{Q}$, it is shown that all primary characteristic classes, i.e. $H^*(BG; k)$, naturally inject into certain Leibniz cohomology groups via an explicit chain map. Moreover, if $f : A \rightarrow B$ is a homomorphism of algebras or rings, the relative Leibniz homology groups $HL_*(f)$ are defined, and if in addition f is surjective with nilpotent kernel, A and B algebras over $\mathbf{Q}$, then there is a natural surjection

$$HL_{*+1}(gl(f)) \rightarrow HC_*(f),$$

where $HC_*(f)$ denotes relative cyclic homology, and $gl(f) : gl(A) \rightarrow gl(B)$ is the induced map on matrices. Here again, the above surjection is realized via an explicit chain map, and offers a relation between Leibniz homology and K-theory, since by work of T. Goodwillie [3], there is an isomorphism

$$K_{*+1}(f) \otimes \mathbf{Q} \rightarrow HC_*(f)$$

between the relative theories when f satisfies the above hypotheses.

Both explicit chain maps mentioned above involve an initial homomorphism

$$\varphi_* : HL_{*+1}(A) \rightarrow HH_*(A)$$

*Supported by the National Science Foundation, grant no. DMS-9704891.

from Leibniz to Hochschild homology, which on the chain level is simply a version of the antisymmetrization map. In many cases, φ_* is seen to be surjective, and from $HH_*(A)$ there are known maps to other types of homologies, such as cyclic homology, or when $A = k[G]$, $HH_*(A)$ maps to $H_*(BG)$. The various homologies can be assembled into a curious commutative diagram:

$$\begin{array}{ccc} HL_{*+1}(A) & \longrightarrow & HH_*(A) \\ \downarrow & & \downarrow \\ H_{*+1}^{\text{Lie}}(A) & \longrightarrow & HC_*(A), \end{array}$$

where $H_*^{\text{Lie}}(A)$ denotes Lie-algebra homology.

The map from Leibniz to Hochschild homology is studied in §2, while the map to $H_*(BG)$ appears in §3, and the map to cyclic homology in the relative case is in the final paragraph. The appendix offers an alternative calculation of the primitives of $HL_*(gl(A))$, and fills a gap in a proof in [8]. Combined with the author's previous results on foliations, Leibniz cohomology contains primary characteristic classes (this paper), secondary characteristic classes [9], and variations of secondary classes [10]. In this way, the HL_* theory can be viewed as a “proto-homology.”

2 From Leibniz to Hochschild Homology

Recall that J.-L. Loday has defined Leibniz homology for the category of Lie algebras, and more generally for the category of Leibniz algebras [7]. In this paper we begin with an associative algebra A over a commutative ring k , and consider A as a Lie algebra via its algebra structure:

$$[a, b] = ab - ba, \quad a, b \in A.$$

In most cases A will be unital, although that assumption is not necessary for the definition of $HL_*(A)$, the Leibniz homology of A . The latter is defined as the homology of the chain complex $CL_*(A)$:

$$k \xleftarrow{0} A \xleftarrow{[\cdot, \cdot]} A^{\otimes 2} \xleftarrow{\quad} \dots \xleftarrow{\quad} A^{\otimes n} \xleftarrow{d} A^{\otimes(n+1)} \xleftarrow{\quad} \dots,$$

where

$$\begin{aligned} d(a_0 \otimes a_1 \otimes \dots \otimes a_n) = \\ \sum_{0 \leq i < j \leq n} (-1)^{j+1} (a_0, a_1, \dots, a_{i-1}, [a_i, a_j], a_{i+1}, \dots, \hat{a}_j, \dots, a_n). \end{aligned} \quad (2.1)$$

The projection from the tensor powers of A to the exterior powers

$$A^{\otimes n} \rightarrow A^{\wedge n}$$

induces a natural map

$$HL_*(A) \rightarrow H_*^{\text{Lie}}(A)$$

from Leibniz homology to Lie-algebra homology, where again the algebra A is viewed as a Lie algebra. We now define a natural map

$$HL_{*+1}(A) \rightarrow HH_*(A)$$

to Hochschild homology, $HH_*(A)$, which when k is a characteristic zero field, yields a commutative diagram:

$$\begin{array}{ccc} HL_{*+1}(A) & \longrightarrow & HH_*(A) \\ \downarrow & & \downarrow \\ H_{*+1}^{\text{Lie}}(A) & \longrightarrow & HC_*(A), \end{array} \quad (2.2)$$

with $HC_*(A)$ denoting cyclic homology.

Recall that $HH_*(A)$ is the homology of the chain complex $CHH_*(A)$:

$$A \xleftarrow{b} A^{\otimes 2} \longleftarrow \dots \longleftarrow A^{\otimes n} \xleftarrow{b} A^{\otimes(n+1)} \longleftarrow \dots,$$

where $b : A^{\otimes(n+1)} \rightarrow A^n$ is defined using the face maps of the cyclic bar construction [6, 1.1.1]

$$\begin{aligned} b &= \sum_{i=0}^n (-1)^i d_i \\ d_i(a_0, a_1, \dots, a_n) &= (a_0, a_1, \dots, a_i a_{i+1}, \dots, a_n), \quad 0 \leq i < n \\ d_n(a_0, a_1, \dots, a_n) &= (a_n a_0, a_1, a_2, \dots, a_{n-1}). \end{aligned} \quad (2.3)$$

Note that $b : A^{\otimes 2} \rightarrow A$ is simply the bracket $[\ , \] : A^{\otimes 2} \rightarrow A$, since

$$b(a_0, a_1) = a_0 a_1 - a_1 a_0,$$

from which follows

$$HL_1(A) \simeq HH_0(A) \simeq HC_0(A) \simeq H_1^{\text{Lie}}(A).$$

In fact, every arrow in diagram (2.2) is an isomorphism in the special case $*$ = 0.

Consider now the map of chain complexes

$$\begin{aligned}\varphi : CL_{*+1}(A) &\rightarrow CHH_*(A) \\ \varphi_n : A^{\otimes(n+1)} &\rightarrow A^{\otimes(n+1)}\end{aligned}\tag{2.4}$$

given by $\varphi_0 = \mathbf{1}$, $\varphi_1 = \mathbf{1}$, and for $n \geq 2$,

$$\begin{aligned}\varphi_n(a_0, a_1, \dots, a_n) &= \\ \sum_{\sigma \in S_n} (\text{sgn } \sigma) (a_0, a_{\sigma^{-1}(1)}, a_{\sigma^{-1}(2)}, \dots, a_{\sigma^{-1}(n)}),\end{aligned}\tag{2.5}$$

where S_n is the symmetric group on n letters. Of course, formula (2.5) could be used to define φ_1 , since S_1 is the trivial group. Also, the summation remains invariant if σ^{-1} is replaced with σ in all subscripts, which reconciles various descriptions of antisymmetrization maps in the literature.

Lemma 2.1. *The k -module homomorphism*

$$\varphi : CL_{*+1}(A) \rightarrow CHH_*(A)$$

is a map of chain complexes.

Proof. It follows at once that $b\varphi_1 = \varphi_0d$. For $n \geq 2$,

$$\begin{aligned}b \circ \varphi_n(a_0, a_1, a_2, \dots, a_n) &= \\ \sum_{\sigma \in S_n} (\text{sgn } \sigma) &\left((a_0 a_{\sigma^{-1}(1)}, a_{\sigma^{-1}(2)}, \dots, a_{\sigma^{-1}(n)}) \right. \\ &- (a_0, a_{\sigma^{-1}(1)} a_{\sigma^{-1}(2)}, a_{\sigma^{-1}(3)}, \dots, a_{\sigma^{-1}(n)}) \\ &+ \dots + (-1)^i (a_0, \dots, a_{\sigma^{-1}(i)} a_{\sigma^{-1}(i+1)}, \dots, a_{\sigma^{-1}(n)}) \\ &\left. + \dots + (-1)^n (a_{\sigma^{-1}(n)} a_0, a_{\sigma^{-1}(1)}, \dots, a_{\sigma^{-1}(n-1)}) \right).\end{aligned}\tag{2.6}$$

On the other hand,

$$\begin{aligned}\varphi_{n-1} \circ d(a_0, a_1, \dots, a_n) &= \\ \varphi_{n-1} \left(\sum_{j=1}^n (-1)^{j+1} ([a_0, a_j], a_1, \dots, \hat{a}_j, \dots, a_n) \right) & \\ + \varphi_{n-1} \left(\sum_{1 \leq i < j \leq n} (-1)^{j+1} (a_0, a_1, \dots, [a_i, a_j], \dots, \hat{a}_j, \dots, a_n) \right).\end{aligned}\tag{2.7}$$

Note that the terms

$$\begin{aligned} \sum_{\sigma \in S_n} (\operatorname{sgn} \sigma) & \left((a_0 a_{\sigma^{-1}(1)}, a_{\sigma^{-1}(2)}, \dots, a_{\sigma^{-1}(n)}) \right. \\ & \left. + (-1)^n (a_{\sigma^{-1}(n)} a_0, a_{\sigma^{-1}(1)}, \dots, a_{\sigma^{-1}(n-1)}) \right) \end{aligned} \quad (2.8)$$

match the terms

$$\varphi_{n-1} \left(\sum_{j=1}^n (-1)^{j+1} ([a_0, a_j], a_1, \dots, \hat{a}_j, \dots, a_n) \right),$$

and the remaining terms in the sums for $b \circ \varphi_n$ and $\varphi_{n-1} \circ d$ also match. Thus,

$$b \circ \varphi_n = \varphi_{n-1} \circ d.$$

□

Corollary 2.2. *There is a natural induced homomorphism*

$$\varphi_* : HL_{*+1}(A) \rightarrow HH_*(A).$$

The chain map φ belongs to the genre of constructions known as antisymmetrization maps [6, 1.3.4]. In fact, it follows from [6, 1.3.5] that φ descends to the exterior powers on the domain

$$\epsilon_n : A \otimes \Lambda^n(A) \rightarrow A^{\otimes(n+1)},$$

and there is an induced map

$$\epsilon_* : H_*^{\operatorname{Lie}}(A; A) \rightarrow HH_*(A) \quad (2.9)$$

where $H_*^{\operatorname{Lie}}(A; A)$ denotes Lie-algebra homology with coefficients in the adjoint representation:

$$\operatorname{ad}(a)(b) = [a, b], \quad a, b \in A.$$

For completeness, recall that $H_*^{\operatorname{Lie}}(A; A)$ is the homology of the complex

$$\begin{aligned} A \xleftarrow{[\cdot, \cdot]} A \otimes A \xleftarrow{\quad} A \otimes A^{\wedge 2} \xleftarrow{\quad} \dots A \otimes A^{\wedge(n-1)} \xleftarrow{d} A \otimes A^{\wedge n} \xleftarrow{\quad} \dots, \\ d(a_0 \otimes a_1 \wedge a_2 \wedge \dots \wedge a_n) = \\ \sum_{1 \leq i < j \leq n} (-1)^{j+1} (a_0 \otimes a_1 \wedge \dots \wedge a_{i-1} \wedge [a_i, a_j] \wedge a_{i+1} \wedge \dots \wedge \hat{a}_j \wedge \dots \wedge a_n) \\ + \sum_{j=1}^n (-1)^{j+1} (\operatorname{ad}(a_0)(a_j) \otimes a_1 \wedge a_2 \wedge \dots \wedge \hat{a}_j \wedge \dots \wedge a_n). \end{aligned}$$

The projection $A^{\otimes(n+1)} \rightarrow A \otimes A^{\wedge n}$ induces a map on homology

$$HL_{*+1}(A) \rightarrow H_*^{\text{Lie}}(A; A). \quad (2.10)$$

Lemma 2.3. *There is a commutative diagram*

$$\begin{array}{ccc} HL_{*+1}(A) & & \\ \downarrow & \searrow \varphi_* & \\ H_*^{\text{Lie}}(A; A) & \xrightarrow{\epsilon_*} & HH_*(A) \end{array}$$

Proof. This follow from corollary (2.2) and maps (2.9) and (2.10) $\square$

When k is a characteristic zero field, the cyclic homology of A may be computed from the complex $C_*^\lambda(A)$ [6, 2.1.4]:

$$A \xleftarrow{b} A^{\otimes 2}/(1-t) \leftarrow \dots A^{\otimes n}/(1-t) \xleftarrow{b} A^{\otimes(n+1)}/(1-t) \leftarrow \dots,$$

where $\mathbf{Z}/(n+1)$ acts on $A^{\otimes(n+1)}$ via

$$t(a_0, a_1, \dots, a_n) = (-1)^n(a_n, a_0, a_1, \dots, a_{n-1}).$$

There is a chain map [6, 10.2.3]

$$\begin{aligned} \theta : \Lambda^{*+1}(A) &\rightarrow C_*^\lambda(A) \\ \theta : \Lambda^{n+1}(A) &\rightarrow A^{\otimes(n+1)}/(1-t) \\ \theta(a_0 \wedge a_1 \wedge \dots \wedge a_n) &= \sum_{\sigma \in S_n} (\text{sgn } \sigma)(a_0, a_{\sigma^{-1}(1)}, a_{\sigma^{-1}(2)} \dots, a_{\sigma^{-1}(n)}), \end{aligned} \quad (2.11)$$

which induces a homomorphism

$$\theta_* : H_{*+1}^{\text{Lie}}(A) \rightarrow HC_*(A).$$

Recall that there is a natural map $I : HH_*(A) \rightarrow HC_*(A)$ [6, 2.2.1], which in the characteristic zero case is induced by the projection

$$\begin{aligned} CHH_*(A) &\rightarrow C_*^\lambda(A) \\ A^{\otimes(n+1)} &\rightarrow A^{\otimes(n+1)}/(1-t). \end{aligned} \quad (2.12)$$

Lemma 2.4. *When k is a characteristic zero field, there is a commutative diagram*

$$\begin{array}{ccc} HL_{*+1}(A) & \xrightarrow{\varphi_*} & HH_*(A) \\ \pi \downarrow & & \downarrow I \\ H_{*+1}^{\text{Lie}}(A) & \xrightarrow{\theta_*} & HC_*(A). \end{array}$$

Proof. This follows immediately, since both φ_* and θ_* are induced by anti-symmetrization maps, and both π and I are induced by projections. $\square$

Of course, when $* = 0$, every arrow in lemma (2.4) is an isomorphism.

The map $\varphi_* : HL_{*+1}(A) \rightarrow HH_*(A)$ has a nice interpretation when A is a smooth algebra over a Noetherian ring k . By definition of smooth, A is assumed to be commutative and unital over k (See [6, 3.4.1] for a discussion of smooth algebras.) Let $\Omega_{A|k}^*$ be the A -module of differential forms, where

$$\Omega_{A|k}^0 = A, \quad \Omega_{A|k}^n = \Lambda_A^n(\Omega_{A|k}^1),$$

and $\Omega_{A|k}^1$ form the Kähler differentials, generated as an A -module by the symbols da , $a \in A$, subject to the relations

$$\begin{aligned} d(\lambda a + \mu b) &= \lambda da + \mu db, \quad \lambda, \mu \in k, \quad a, b \in A \\ d(ab) &= a(db) + b(da), \quad a, b \in A \end{aligned} \tag{2.13}$$

By a theorem of Hochschild, Kostant, and Rosenberg [4], when A is smooth over k , the antisymmetrization

$$\epsilon_* : \Omega_{A|k}^* \rightarrow HH_*(A) \tag{2.14}$$

is an isomorphism of graded algebras. Of course, for any commutative algebra A , $[a, b] = 0$ for all $a, b \in A$, and

$$HL_*(A) \simeq T(A), \tag{2.15}$$

where $T(A) = \sum_{k \geq 0} A^{\otimes k}$ denotes the tensor algebra on A .

Lemma 2.5. *If A is smooth over k , then the natural map*

$$\varphi_* : HL_{*+1}(A) \rightarrow HH_*(A)$$

is surjective.

Proof. There is a surjective k -module homomorphism

$$HL_{n+1}(A) \simeq A^{\otimes(n+1)} \rightarrow \Omega_{A|k}^n$$

given on homogeneous elements by

$$p(a_0 \otimes a_1 \otimes \dots \otimes a_n) = a_0 da_1 \wedge da_2 \wedge \dots \wedge da_n.$$

The result now follows from the commutativity of the diagram

$$\begin{array}{ccc} HL_{*+1}(A) & & \\ p \downarrow & \searrow \varphi_* & \\ \Omega_{A|k}^* & \xrightarrow[\epsilon_*]{} & HH_*(A). \end{array}$$

□

When A is the smooth algebra $C^\infty(M)$ of real-valued differentiable functions on a differentiable manifold M , the reader is invited to identify the various homology groups in lemma (2.4) using the calculations of $HC_*(C^\infty(M))$ in terms of de Rham cohomology [6, 3.4.12].

In general, the map $\varphi_* : HL_{*+1}(A) \rightarrow HH_*(A)$ is not surjective, even in the commutative case, since from the definition of φ

$$\text{Im} [\varphi_*(HL_{n+1}(A))] \subset HH_n^{(n)}(A),$$

where $HH_n^{(n)}(A)$ denotes the n -th summand in the λ -decomposition of Hochschild homology over $\mathbf{Q}$. See [6, 4.5] for details about this decomposition. Of course, $HH_n^{(n)}(A)$ is not necessarily isomorphic to $HH_n(A)$. In the next section we investigate another important case in which φ_* is surjective.

3 Relation to Characteristic Classes

In this paragraph we prove that the natural map

$$\varphi_* : HL_{*+1}(gl(A)) \rightarrow HH_*(gl(A)) \simeq HH_*(A)$$

is onto for a unital algebra over a characteristic zero field k . Furthermore, when A is the group ring $k[G]$, there is a surjective homomorphism

$$HH_*(k[G]) \rightarrow H_*(k[G]) := H_*(BG; k),$$

and, in fact, $H_*(BG; k)$ is a direct summand of $HH_*(k[G])$. Thus, on cohomology

$$H^*(BG; k) \rightarrow HL^{*+1}(gl(A))$$

is injective, and $HL^{*+1}(gl(A))$ contains all characteristic classes in

$$H^*(BG; k).$$

Specifically, let

$$gl(A) = \varinjlim_n gl_n(A)$$

be the Lie algebra of infinite matrices over A with finitely many nonzero entries. Note that the Lie algebra structure on $gl(A)$ is actually induced from the ring structure of

$$M(A) = \varinjlim_n M_n(A),$$

where $M_n(A) = gl_n(A)$ is the ring of all $n \times n$ matrices over A . Recall that [2] [6, 10.6.5]

$$HL_*(gl(A)) \simeq T(HH_*(A)[1]), \quad (3.1)$$

where T denotes the tensor algebra. See also [8]. To understand the map φ_* on homology, the isomorphism in equation (3.1) must be understood on the chain level. The chain complex $CL_*(gl(A))$ is quasi-isomorphic to $L_*(A)$ [6, 10.6.7]:

$$k[S_1] \otimes_k A \longleftarrow k[S_2] \otimes_k A^{\otimes 2} \longleftarrow \dots \longleftarrow k[S_n] \otimes_k A^{\otimes n} \longleftarrow \dots, \quad (3.2)$$

where S_n denotes the symmetric group on n letters. There is an explicit chain map

$$\begin{aligned} CL_*(gl(A)) &\rightarrow L_*(A) \\ [gl(A)]^{\otimes n} &\rightarrow ([gl(A)]^{\otimes n})_{gl(k)} \xrightarrow{\Theta} k[S_n] \otimes A^{\otimes n} \end{aligned} \quad (3.3)$$

with $[gl(A)]^{\otimes n} \rightarrow ([gl(A)]^{\otimes n})_{gl(k)}$ being the projection onto the quotient by the adjoint action. For $\sigma \in S_n$ and E_{ij}^a the elementary matrix with only one possible nonzero entry $a \in A$ in the ij position,

$$\Theta(E_{1\sigma(1)}^{a_1} \otimes E_{2\sigma(2)}^{a_2} \otimes \dots \otimes E_{n\sigma(n)}^{a_n}) = \sigma \otimes (a_1, a_2, \dots, a_n) \quad (3.4)$$

For more details, see [6, 10.2.11].

Now, $HH_*(A)[1]$ is the direct summand of $HL_*(gl(A))$ that arises from the homology of the complex $P_*(A)$:

$$k[U_1] \otimes_k A \longleftarrow k[U_2] \otimes_k A^{\otimes 2} \longleftarrow \dots \longleftarrow k[U_n] \otimes_k A^{\otimes n} \longleftarrow \dots, \quad (3.5)$$

where U_n is the conjugacy class of the cyclic shift in S_n . The complex $P_*(A)$ is a summand of $L_*(A)$ [8], and an alternative calculation of $H_*(P_*(A))$ appears in the appendix. Also needed for the identification of $\text{Im}(\varphi_*)$ is an explicit description of the trace map isomorphism from Morita invariance

$$\text{tr}_* : HH_*(gl(A)) \rightarrow HH_*(A).$$

For ${}_p M = [{}_p m_{ij}] \in gl(A)$,

$$\begin{aligned} \text{tr}({}_0 M \otimes {}_1 M \otimes \dots \otimes {}_n M) = \\ \sum {}_0 m_{i_1 i_2} \otimes {}_1 m_{i_2 i_3} \otimes {}_2 m_{i_3 i_4} \otimes \dots \otimes {}_n m_{i_{n+1} i_1}, \end{aligned} \quad (3.6)$$

where the sum is over all indices $i_1, i_2, \dots, i_{n+1}$.

Lemma 3.1. *Let A be a unital algebra over a characteristic zero field k . Then the composition*

$$\text{tr}_* \circ \varphi_* : HL_{*+1}(gl(A)) \rightarrow HH_*(A)$$

is surjective.

Proof. The summand of $HL_{*+1}(gl(A))$ isomorphic to $HH_*(A)[1]$ can be represented via chains which are k -linear combinations of terms:

$$E_{1\tau(1)}^{a_1} \otimes E_{2\tau(2)}^{a_2} \otimes \dots \otimes E_{n\tau(n)}^{a_n},$$

where τ is the cyclic shift given by the cycle $(1, 2, 3, \dots, n)$ [8]. It is enough to compute

$$\begin{aligned} \text{tr} \circ \varphi(E_{1\tau(1)}^{a_1} \otimes E_{2\tau(2)}^{a_2} \otimes \dots \otimes E_{n\tau(n)}^{a_n}) \\ = \text{tr} \circ \varphi(E_{12}^{a_1} \otimes E_{23}^{a_2} \otimes \dots \otimes E_{n1}^{a_n}) \\ = a_1 \otimes a_2 \otimes \dots \otimes a_n. \end{aligned}$$

Thus, any element in $HH_*(A)$ can be represented as a chain in $CL_{*+1}(gl(A))$. $\square$

Consider the case of a group ring $A = k[G]$, and the associated bar construction $\{G^n\}_{n \geq 0}$. The face maps $d_i : G^n \rightarrow G^{n-1}$ are given by

$$d_i(g_1, g_2, \dots, g_n) = \begin{cases} (g_2, g_3, \dots, g_n) & i = 0, \\ (g_1, \dots, g_i g_{i+1}, \dots, g_n) & 1 \leq i \leq n-1, \\ (g_1, g_2, \dots, g_{n-1}) & i = n, \end{cases}$$

with $H_*(BG)$ denoting the homology of the complex

$$B_*(G) = k[G^n] \simeq (k[G])^{\otimes n}, \quad n \geq 0.$$

There are natural simplicial maps

$$\begin{aligned} \pi : CHH_*(k[G]) &\rightarrow B_*(G) \\ \pi : k[G^{n+1}] &\rightarrow k[G^n] \\ \pi(g_0, g_1, g_2, \dots, g_n) &= (g_1, g_2, \dots, g_n) \\ \iota : B_*(G) &\rightarrow CHH_*(k[G]) \\ \iota : k[G^n] &\rightarrow k[G^{n+1}] \\ \iota(g_1, g_2, \dots, g_n) &= ((g_1 g_2 \cdots g_n)^{-1}, g_1, g_2, \dots, g_n), \end{aligned}$$

and the composition $\pi \circ \iota$ is the identity on $B_*(G)$. Thus follows the known lemma [1]

Lemma 3.2. [1] *The graded group $H_*(BG)$ is a direct summand of*

$$HH_*(k[G])$$

for any coefficient ring k .

By composing the chain maps π , tr , and φ , we have

Theorem 3.3. *Let k be a characteristic zero field and $A = k[G]$. Then*

$$(\pi \circ \text{tr} \circ \varphi)_* : HL_{*+1}(gl(A)) \rightarrow H_*(BG; k)$$

is surjective, and

$$(\pi \circ \text{tr} \circ \varphi)^* : H^*(BG; k) \rightarrow HL^{*+1}(gl(A))$$

is injective.

Thus, all characteristic classes over k naturally inject into Leibniz cohomology.

4 Relation to K-theory

A fundamental theorem of T. Goodwillie [3] relates relative algebraic K-theory, $K_*(f)$, to relative cyclic homology, $HC_*(f)$. In particular, if $f : R \rightarrow S$ is a homomorphism of simplicial rings such that the induced map $\pi_0(R) \rightarrow \pi_0(S)$ is surjective with nilpotent kernel, then

$$K_n(f) \otimes \mathbf{Q} \simeq HC_{n-1}(f) \otimes \mathbf{Q}. \quad (4.1)$$

Although for Leibniz homology we are working in the category of discrete rings, the following results may be extended to the simplicial case. For a homomorphism $f : A \rightarrow B$ of discrete rings, we define the relative Leibniz homology groups, $HL_*(f)$, we then consider the map on matrices $gl(f) : gl(A) \rightarrow gl(B)$, and show that over a characteristic zero field, the composition

$$(\mathrm{tr} \circ \varphi)_* : HL_{*+1}(gl(f)) \rightarrow HH_*(f)$$

is onto. If furthermore, $f : A \rightarrow B$ is surjective with nilpotent kernel, then

$$I : HH_*(f) \rightarrow HC_*(f)$$

is also surjective, as well as the composition

$$I \circ (\mathrm{tr} \circ \varphi)_* : HL_{*+1}(gl(f)) \rightarrow HC_*(f). \quad (4.2)$$

Above, all relative homology groups are taken with coefficients in a characteristic zero field.

Recall briefly the construction of relative homology in a general setting [11, p. 46–47]. Let $f : C_n \rightarrow C'_n$, $n \geq 0$, be a homomorphism between any two chain complexes C_* and C'_* , and define the mapping cone of f as

$$M_n(f) = C_{n-1} \oplus C'_n, \quad n \geq 1,$$

with boundary map

$$\partial(c, c') = (-\partial c, \partial c' + fc)$$

By definition, $H_*(f)$ are the homology groups of $M_*[-1] = M_{*+1}$, which fit into a long exact sequence

$$\begin{aligned} \dots &\longrightarrow H_n(C_*) \xrightarrow{f_*} H_n(C'_*) \xrightarrow{\alpha_*} H_n(f) \xrightarrow{p_*} H_{n-1}(C_*) \longrightarrow \dots \\ \dots &\longrightarrow H_1(f) \xrightarrow{p_*} H_0(C) \xrightarrow{f_*} H_0(C'_*) \xrightarrow{\alpha_*} H_0(f) \longrightarrow 0, \end{aligned} \quad (4.3)$$

where $p : M_*(f) \rightarrow C_{*-1}$ and $\alpha : C'_* \rightarrow M_*(f)$ are given by

$$p((c, c')) = c, \quad \alpha(c') = (0, c').$$

If $f : A \rightarrow B$ is a homomorphism of discrete rings, then the above construction yields $HL_*(f)$ by considering the chain map $f : CL_*(A) \rightarrow CL_*(B)$. Moreover, the relative homology group construction is functorial, and there are natural maps

$$\begin{aligned} \varphi_* : HL_{*+1}(f) &\rightarrow HH_*(f) \\ \text{tr}_* : HH_*(gl(f)) &\rightarrow HH_*(f) \\ I : HH_*(f) &\rightarrow HC_*(f). \end{aligned}$$

Theorem 4.1. *Suppose that A and B are unital algebras over a characteristic zero field k , and $f : A \rightarrow B$ is an algebra homomorphism. For the map $gl(f) : gl(A) \rightarrow gl(B)$, the composition*

$$(\text{tr} \circ \varphi)_* : HL_{*+1}(gl(f)) \rightarrow HH_*(gl(f)) \rightarrow HH_*(f)$$

is surjective.

Proof. From equation (3.1)

$$\begin{aligned} HL_*(gl(A)) &\simeq T(HH_*(A)[1]) \\ HL_*(gl(B)) &\simeq T(HH_*(B)[1]) \end{aligned}$$

From equation (3.5) the summand of $HL_*(gl(A))$ isomorphic to $HH_*(A)[1]$ can be computed from the complex $P_*(A)$, and similarly for $gl(B)$. Consider the k -linear homomorphism

$$\begin{aligned} P(f) : P_*(A) &\rightarrow P_*(B) \\ P(f) : k[U_n] \otimes A^{\otimes n} &\rightarrow k[U_n] \otimes B^{\otimes n} \\ P(f)(\sigma \otimes (a_1, a_2, \dots, a_n)) &= \sigma \otimes (f(a_1), f(a_2), \dots, f(a_n)). \end{aligned}$$

We have the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} \longrightarrow & H_{n+1}(P_*(A)) & \longrightarrow & H_{n+1}(P_*(B)) & \xrightarrow{\alpha_*} & H_{n+1}(P(f)) & \longrightarrow \\ & \downarrow & & \downarrow & & \downarrow & \\ \longrightarrow & HH_n(A) & \longrightarrow & HH_n(B) & \xrightarrow{\alpha_*} & HH_n(f) & \longrightarrow, \end{array}$$

where the dimensions in the top row are inherited from the chain complex for Leibniz homology. By the 5-lemma, $H_*(P(f)) \simeq HH_*(f)[1]$. Let z denote an element in the mapping cone

$$CL_*(gl(A)) \rightarrow CL_*(gl(B))$$

of the form

$$(E_{1\tau(1)}^{a_1} \otimes E_{2\tau(2)}^{a_2} \otimes \dots \otimes E_{n\tau(n)}^{a_n}, E_{1\tau'(1)}^{b_1} \otimes E_{2\tau'(2)}^{b_2} \otimes \dots \otimes E_{n+1,\tau'(n+1)}^{b_{n+1}}),$$

where τ is the cyclic shift in S_n and τ' is the cyclic shift in S_{n+1} . The chain map Θ in equation (3.4) may be defined on the respective mapping cones, and

$$\Theta(z) = (\tau \otimes (a_1, a_2, \dots, a_n), \tau' \otimes (b_1, b_2, \dots, b_{n+1})),$$

where the latter is in fact an element of $M(P(f))$. Also, at the level of mapping cones, we have

$$(\text{tr} \circ \varphi)(z) = ((a_1, a_2, \dots, a_n), (b_1, b_2, \dots, b_{n+1})).$$

The theorem follows, since the isomorphism

$$H_*(P(f)) \simeq HH_*(f)[1]$$

is realized by sending the class of $\Theta(z)$ to the class of $(\text{tr} \circ \varphi)(z)$. $\square$

For relative cyclic homology, there is a long exact sequence

$$\dots \longrightarrow HH_n(f) \xrightarrow{I} HC_n(f) \xrightarrow{S} HC_{n-2}(f) \xrightarrow{B} HH_{n-1}(f) \xrightarrow{I} \dots$$

By work of Goodwillie, we have

Lemma 4.2. *If $f : A \rightarrow B$ is a homomorphism of unital algebras over a characteristic zero field with nilpotent kernel, then the map*

$$I : HH_n(f) \rightarrow HC_n(f)$$

is surjective.

Proof. From [3, p. 399], the map $S : HC_n(f) \rightarrow HC_{n-2}(f)$ is zero. $\square$

Corollary 4.3. *Under the hypotheses of lemma (4.2), the natural map*

$$I \circ (\text{tr} \circ \varphi)_* : HL_{*+1}(gl(f)) \rightarrow HC_*(f)$$

is surjective.

5 Appendix: The Homology of $P_*(A)$

The results of this paper concerning the surjectivity of the homomorphism

$$HL_{*+1}(gl(A)) \rightarrow HH_*(A)$$

rely heavily on the calculation of the homology of the complex $P_*(A)$:

$$k[U_1] \otimes_k A \longleftarrow k[U_2] \otimes_k A^{\otimes 2} \longleftarrow \dots \longleftarrow k[U_{n+1}] \otimes_k A^{\otimes(n+1)} \longleftarrow \dots,$$

which form the primitive elements of $HL_*(gl(A))$. In this appendix we offer a calculation of $H_*(P_*(A))$ which fills a gap in a previous proof, namely lemma (2.6) of [8]. We prove that if A is a unital k -algebra, then

$$H_*(P_*(A)) \simeq HH_*(A),$$

where now $P_n(A) = k[U_{n+1}] \otimes_k A^{\otimes(n+1)}$.

Recall that U_n is the conjugacy class of the cyclic shift in the symmetric group S_n . The collection $\{U_{n+1}\}_{n \geq 0}$ form a presimplicial set with face maps d_i [8], but lack degeneracies. Let $N^{\text{cy}}(A)$ be the cyclic bar construction on A with faces given in equation (2.3). Let

$$k[U_{*+1}] \otimes N^{\text{cy}}(A)$$

be the presimplicial k -module with

$$\{k[U_{*+1}] \otimes N^{\text{cy}}(A)\}_n = k[U_{n+1}] \otimes A^{\otimes(n+1)},$$

and face maps

$$d_i(\sigma \otimes \vec{a}) = d_i(\sigma) \otimes d_i(\vec{a}).$$

The complex $P_*(A)$ is simply $k[U_{*+1}] \otimes N^{\text{cy}}(A)$ together with its boundary map constructed as the alternating sum of the face maps. In [8] it is shown that the complex $k[U_{*+1}]$ is acyclic, but without degeneracies, the Eilenberg-Zilber and Künneth theorems cannot be applied to calculate $H_*(P_*(A))$. To remedy this, we invoke the Dold-Kan functor [5], which to any chain complex K_* (over k), associates the simplicial k -module

$$D_n(K_*) = \text{Hom}(C(\Delta[n]), K_*),$$

where Hom denotes chain maps over k . Also, $\Delta[n]$ is the simplicial model for the n -simplex, $C'(\Delta[n])$ is the free k -module on the elements of $\Delta[n]$, and

$$C(\Delta[n]) = C'(\Delta[n]) / {}^e C(\Delta[n]),$$

with ${}^e C(\Delta[n])$ denoting the submodule of $C'(\Delta[n])$ generated by the degenerate elements. See [5] for further details.

Using properties of the Dold-Kan functor, the complexes

$$k[U_{*+1}] \otimes N^{\text{cy}}(A), \quad D_*(k[U_{*+1}]) \otimes N^{\text{cy}}(A)$$

are quasi-isomorphic, as well as the complexes

$$k[U_{*+1}] \otimes N^{\text{cy}}(A), \quad D_*(k[U_{*+1}] \otimes N^{\text{cy}}(A)).$$

Define a chain map

$$\begin{aligned} \psi : D_*(k[U_{*+1}]) \otimes N^{\text{cy}}(A) &\rightarrow D_*(k[U_{*+1}] \otimes N^{\text{cy}}(A)) \\ \psi : \text{Hom}(C(\Delta[n]), k[U_{*+1}]) \otimes A^{\otimes(n+1)} &\rightarrow \text{Hom}(C(\Delta[n]), k[U_{*+1}] \otimes A^{**+1}) \end{aligned}$$

as follows. Recall that $\delta^i : [n-1] \rightarrow [n]$, $i = 0, 1, 2, \dots, n$ is given by

$$\delta^i(j) = \begin{cases} j, & j < i \\ j+1, & j \geq i. \end{cases}$$

Let $\alpha \in (\Delta[n])_q$, $\alpha : [q] \rightarrow [n]$, $q \leq n$. Then

$$\begin{aligned} \alpha &= \delta^{i_{n-1}} \circ \dots \circ \delta^{i_2} \circ \delta^{i_1} \\ [q] &\xrightarrow{\delta^{i_1}} [q+1] \xrightarrow{\delta^{i_2}} [q+2] \longrightarrow \dots \longrightarrow [n-1] \xrightarrow{\delta^{i_{n-q}}} [n], \end{aligned}$$

and the factorization of α is well-defined up to cosimplicial identities. Suppose that $\vec{a} \in A^{\otimes(n+1)}$ and

$$g : C(\Delta[n]) \rightarrow k[U_{*+1}]$$

is a chain map. We define

$$\psi(g \otimes \vec{a})(\alpha) = g(\alpha) \otimes (d_{i_1} \circ d_{i_2} \circ \dots \circ d_{i_{n-q}}(\vec{a})).$$

Now, $H_*(D_*(k[u_{*+1}]) \otimes N^{\text{cy}}(A)) \simeq HH_*(A)$, since $D_*(k[U_{*+1}])$ is an acyclic simplicial k -module. Also,

$$H_*(D_*(k[u_{*+1}] \otimes N^{\text{cy}}(A))) \simeq H_*(k[U_{*+1}] \otimes N^{\text{cy}}(A)).$$

Define an inclusion of chain complexes

$$\begin{aligned} N^{\text{cy}}(A) &\rightarrow k[U_{*+1}] \otimes N^{\text{cy}}(A) \\ A^{\otimes(n+1)} &\rightarrow k[U_{n+1}] \otimes A^{\otimes(n+1)} \\ \vec{a} &\mapsto \tau_{n+1} \otimes \vec{a}, \end{aligned}$$

where τ_{n+1} is the cyclic shift in U_{n+1} . We then have a commutative diagram of chain complexes, where the diagonal and vertical arrows are inclusions:

$$\begin{array}{ccc} N^{\text{cy}}(A) & \longrightarrow & k[U_{*+1}] \otimes N^{\text{cy}}(A) \\ \downarrow & \swarrow & \downarrow \\ D_*(k[U_{*+1}]) \otimes N^{\text{cy}}(A) & \xrightarrow[\psi]{} & D_*(k[U_{*+1}] \otimes N^{\text{cy}}(A)). \end{array}$$

On homology

$$\begin{array}{ccc} HH_*(A) & \longrightarrow & H_*(k[U_{*+1}] \otimes N^{\text{cy}}(A)) \\ \simeq \downarrow & \swarrow & \downarrow \simeq \\ H_*(D_*(k[U_{*+1}]) \otimes N^{\text{cy}}(A)) & \xrightarrow[\psi_*]{} & H_*(D_*(k[U_{*+1}] \otimes N^{\text{cy}}(A))). \end{array} \quad (5.1)$$

It follows that

$$HH_*(A) \rightarrow H_*(k[U_{*+1}] \otimes N^{\text{cy}}(A))$$

is an isomorphism, and $HH_*(A) \simeq H_*(P_*(A))$. In fact, every arrow in diagram (5.1) is an isomorphism.

References

- [1] Burghilea, D., “The Cyclic Homology of the Group Rings,” *Comment. Math. Helv.* 60, (1985), 354–365.
- [2] Cuvier, C., “Homologie de Leibniz et homologie de Hochschild,” *Comptes Rendus Acad. Sci. Paris Sér. A–B*, 313, (1991), 569–572.

- [3] Goodwillie, T., “Relative algebraic K-theory and cyclic homology,” *Annals of Math.*, 124, (1986), 347–402.
- [4] Hochschild, G., Kostant, B., Rosenberg, A., “Differential Forms on Regular Affine Algebras,” *Trans. AMS*, 102, (1962), 383–408.
- [5] Lamotke, K., *Semisimpliziale Algebraische Topologie*, Grund. der Math. Wissen., 147, Springer Verlag, 1968.
- [6] Loday, J.-L., *Cyclic Homology*, second ed., Grund. der Math. Wissen., 301, Springer Verlag, 1998.
- [7] Loday, J.-L., Pirashvili, T., “Universal Enveloping Algebras of Leibniz Algebras and (Co)-homology,” *Math. Annalen*, 296 (1993), 139–158.
- [8] Lodder, J., “Leibniz Homology and the James Model,” *Math. Nachr.*, 175, (1995), 209–229.
- [9] Lodder, J., “Leibniz Cohomology for Differentiable Manifolds,” *Ann. Inst. Fourier*, Grenoble, 48, 1, (1998), 73–95.
- [10] Lodder, J., “Rigidity of Secondary Characteristic Classes,” *Diff. Geom. and its Applications*, 12, (2000), 207–218.
- [11] Mac Lane, S., *Homology*, Grund. der Math. Wissen., 114, Springer Verlag, 1975.